



Restriction estimates via the derivatives of the heat semigroup and connexion with dispersive estimates

Frederic Bernicot, El Maati Ouhabaz

► To cite this version:

Frederic Bernicot, El Maati Ouhabaz. Restriction estimates via the derivatives of the heat semigroup and connexion with dispersive estimates. *Mathematical Research Letters*, 2013, 20 (6), pp.1047-1058. hal-00812285

HAL Id: hal-00812285

<https://hal.science/hal-00812285>

Submitted on 11 Apr 2013

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

RESTRICTION ESTIMATES VIA THE DERIVATIVES OF THE HEAT SEMIGROUP AND CONNEXION WITH DISPERSIVE ESTIMATES

FRÉDÉRIC BERNICOT & EL MAATI OUHABAZ

ABSTRACT. We consider an abstract non-negative self-adjoint operator H on an L^2 -space. We derive a characterization for the restriction estimate $\|dE_H(\lambda)\|_{L^p \rightarrow L^{p'}} \leq C\lambda^{\frac{d}{2}(\frac{1}{p}-\frac{1}{p'})-1}$ in terms of higher order derivatives of the semigroup e^{-tH} . We provide an alternative proof of a result in [1] which asserts that dispersive estimates imply restriction estimates. We also prove $L^p - L^{p'}$ estimates for the derivatives of the spectral resolution of H .

1. INTRODUCTION AND MAIN RESULTS

Let (X, μ) be a measured space. That is X is a non-empty set endowed with a positive measure μ . We consider a non-negative self-adjoint operator H on $L^2 = L^2(X, \mu)$. We denote by dE_H the spectral resolution of H . Since we will be interested in $L^p - L^{p'}$ estimates for $dE_H(\lambda)$ we shall assume throughout this note that the spectrum of H is continuous. The $L^p - L^{p'}$ norm will be denoted by $\|dE_H(\lambda)\|_{L^p \rightarrow L^{p'}}$ and p' is the conjugate number of p .

We first discuss the Euclidean Laplacian. Suppose that $X = \mathbb{R}^d$ and $H = -\Delta$ (the positive Laplace operator) on $L^2(\mathbb{R}^d)$. It is a well-known fact that as a consequence of the Stein-Tomas estimates for the restricted Fourier transform to the unit sphere, the spectral measure $dE_{-\Delta}(\lambda)$ is a bounded operator from L^p into $L^{p'}$ for all $p \leq \frac{2d+2}{d+3}$. In addition,

$$\|dE_{-\Delta}(\lambda)\|_{L^p \rightarrow L^{p'}} \leq C\lambda^{\frac{d}{2}(\frac{1}{p}-\frac{1}{p'})-1}, \quad \lambda > 0.$$

Such estimate is sometimes referred to as the $(p, 2)$ restriction estimate of Stein-Tomas. We refer to the introductions of the papers [3] and [1] for more details about this.

The above restriction estimate was extended to the setting of asymptotically conic manifolds in [3]. In the paper [1] the restriction estimate

$$(1.1) \quad \|dE_H(\lambda)\|_{L^p \rightarrow L^{p'}} \leq C\lambda^{\frac{d}{2}(\frac{1}{p}-\frac{1}{p'})-1},$$

was studied in an abstract setting. (Here d is any positive constant). It is also proved there that (1.1) holds for several operators.

One of the aims of this note is to prove other characterizations of (1.1) in an abstract setting. In the following result we show that the restriction estimate for H can be characterized in terms of higher order derivatives of the corresponding semigroup e^{-tH} . More precisely,

Theorem 1.1. *Let d be a positive constant and fix $p \in [1, 2)$. The following assertions are equivalent.*

Date: April 11, 2013.

2000 Mathematics Subject Classification. 47D03 ; 35P05 ; 42B15.

Key words and phrases. Restriction estimates ; semigroup ; spectral multipliers ; dispersive estimates.

The two authors are partly supported by the ANR under the project "Harmonic Analysis at its Boundaries" no. ANR-12-BS01-0013. The first author is also partly supported by the ANR under the project AFoMEN no. 2011-JS01-001-01.

- 1) The restriction estimate (1.1) holds for every $\lambda > 0$;
 2) There exists a positive constant C such that

$$(1.2) \quad \|H^N e^{-tH}\|_{L^p \rightarrow L^{p'}} \leq C(N-1)! N^{\frac{d}{2}(\frac{1}{p}-\frac{1}{p'})} t^{-N-\frac{d}{2}(\frac{1}{p}-\frac{1}{p'})},$$

for all $t > 0$ and all $N \in \mathbb{N}$;

- 3) There exists a positive constant C such that

$$(1.3) \quad \|F(H)\|_{L^p \rightarrow L^{p'}} \leq C R^{\frac{d}{2}(\frac{1}{p}-\frac{1}{p'})} \int_{\mathbb{R}} |F(s)| \frac{ds}{s},$$

for all $R > 0$ and bounded measurable function F with supported in $[0, R]$.

The main novelty here is the characterization of (1.1) by (1.2). The equivalence of (1.1) and (1.3) is in the spirit of Proposition 2.4, Section 2.2 in [1]. Note however that in contrast to that proposition in [1] we do not assume here that the volume of balls in X is polynomial. Moreover, the L^1 norm in the RHS in (1.3) is taken *w.r.t.* $\frac{ds}{s}$ rather than ds as in [1] and it is obvious that

$$\int_0^R |F(s)| \frac{ds}{s} = \int_0^1 |F(Rs)| \frac{ds}{s} \geq \int_0^1 |F(Rs)| ds.$$

One of the main ingredients in the proof of Theorem 1.1 is the following result which expresses the spectral measure in terms of the semigroup. We denote by $\langle \cdot, \cdot \rangle$ the scalar product of L^2 . We have

Theorem 1.2. *Consider a bounded and uniformly ρ -Hölder function ϕ (for some $\rho \in (0, 1]$). Then for every $f, g \in L^2$ we have*

$$\lim_{N \rightarrow \infty} \frac{1}{(N-1)!} \int_0^\infty \phi(s^{-1}) \langle ((N-1)sH)^N e^{-s(N-1)H} f, g \rangle \frac{ds}{s} = \langle \phi(H) f, g \rangle.$$

A useful consequence of the latter theorem is the following equality for the derivatives of $dE_H(\lambda)$ (in which the limit has to be understood in the weak sense)

$$\frac{d^k}{d\lambda^k} dE_H(\lambda) = \lim_{N \rightarrow \infty} \frac{1}{N!} \frac{d^k}{d\lambda^k} \left[\lambda^{-1} (N\lambda^{-1}H)^{N+1} e^{-N\lambda^{-1}H} \right],$$

for $\lambda > 0$.

As an application we show that dispersive estimates for H imply the restriction estimate (1.1) as well as $L^p - L^{p'}$ estimates for the derivatives $\frac{d^k}{d\lambda^k} dE_H(\lambda)$ for $k \leq d/2 - 1$. The estimates for the derivatives is new whereas the case $k = 0$ was already proved in [1].

We finish this introduction by explaining why it is interesting to prove the restriction estimate (1.1). Let us assume now that (X, μ) is equipped with a metric ρ and assume that for every $x \in X$, $r > 0$, the volume $\mu(B(x, r))$ of the open ball $B(x, r)$ satisfies

$$c_1 r^d \leq \mu(B(x, r)) \leq c_2 r^d,$$

where c_1 and c_2 are positive constants. Suppose in addition that H satisfies the finite speed of propagation property, that is the support of the kernel of $\cos(t\sqrt{H})$ is contained in $\{(x, y) \in X \times X, \rho(x, y) \leq t\}$. Under these assumptions, it is proved in [1] (see also [3] for the first assertion) that the restriction estimate implies sharp spectral multiplier theorems. More precisely,

Theorem 1.3. *Suppose that the restriction estimate (1.1) holds for some fixed $p \in [1, 2)$. Then the following assertions hold.*

- (i) **Compactly supported multipliers:** Let F be an even function with support in $[-1, 1]$ and $F \in W^{\beta, 2}(\mathbb{R})$ for some $\beta > d(1/p - 1/2)$. Then $F(H)$ is bounded on $L^p(X)$, and

$$\sup_{t>0} \|F(tH)\|_{L^p \rightarrow L^p} \leq C \|F\|_{W^{\beta, 2}}.$$

- (ii) **General multipliers:** Suppose that F is an even bounded Borel function which satisfies $\sup_{t>0} \|\eta(\cdot)F(t\cdot)\|_{W^{\beta, 2}} < \infty$ for some $\beta > \max\{d(1/p - 1/2), 1/2\}$ and some non-trivial function $\eta \in C_c^\infty(0, \infty)$. Then $F(H)$ is bounded on $L^r(X)$ for all $p < r < p'$. In addition,

$$\|F(H)\|_{L^r \rightarrow L^r} \leq C_\beta \left(\sup_{t>0} \|\eta(\cdot)F(t\cdot)\|_{W^{\beta, 2}} + |F(0)| \right).$$

A version of this theorem for general doubling spaces is proved in [1]. One can apply Theorem 1.3 to prove summability results for Bochner-Riesz means on L^p -spaces.

2. PROOFS

We start with the proof of Theorem 1.2. We shall write \lesssim for \leq up to a non relevant constant C .

Proof of Theorem 1.2. Let us set $c_N^{-1} := \int_0^\infty x^N e^{-x} \frac{dx}{x} = (N-1)!$. By polarization, it suffices to prove that for every function $f \in L^2$

$$(2.1) \quad \lim_{N \rightarrow \infty} c_N \int_0^\infty \phi(s^{-1}) \langle ((N-1)sH)^N e^{-s(N-1)H} f, f \rangle \frac{ds}{s} = \langle \phi(H)f, f \rangle.$$

We have

$$(2.2) \quad \begin{aligned} c_N \int_0^\infty \phi(s^{-1}) \langle ((N-1)sH)^N e^{-s(N-1)H} f, f \rangle \frac{ds}{s} \\ = c_N \int_0^\infty \int_0^\infty \phi(s^{-1}) ((N-1)s\lambda)^N e^{-(N-1)s\lambda} \langle dE_\lambda f, f \rangle \frac{ds}{s} \\ (2.3) \quad = \int_0^\infty \mu_N(\lambda) \langle dE_\lambda f, f \rangle, \end{aligned}$$

where

$$\mu_N(\lambda) := c_N \int_0^\infty \phi(s^{-1}) ((N-1)s\lambda)^N e^{-(N-1)s\lambda} \frac{ds}{s}.$$

Due to the constant c_N , it is clear that the continuous function μ_N is bounded by $\|\phi\|_{L^\infty}$. Therefore, it is enough to prove that $\mu_N(\lambda)$ converges to the function $\phi(\lambda)$ for all $\lambda > 0$ and then conclude by the dominated convergence theorem.

Taking the difference yields

$$\begin{aligned} |\mu_N(\lambda) - \phi(\lambda)| &= c_N \left| \int_0^\infty [\phi(s^{-1}) - \phi(\lambda)] ((N-1)s\lambda)^N e^{-(N-1)s\lambda} \frac{ds}{s} \right| \\ &\leq \int_0^\infty |\phi(s^{-1}) - \phi(\lambda)| c_N ((N-1)s\lambda)^N e^{-(N-1)s\lambda} \frac{ds}{s}, \end{aligned}$$

Using Stirling's formula

$$c_N = [(N-1)!]^{-1} \simeq \left(\frac{e}{N-1} \right)^{N-1} (2\pi N)^{-\frac{1}{2}}$$

we obtain for large enough N and uniformly with respect to λ

$$|\mu_N(\lambda) - \phi(\lambda)| \lesssim N^{\frac{1}{2}} \int_0^\infty |\phi(s^{-1}) - \phi(\lambda)| (s\lambda)^N e^{-(N-1)(s\lambda-1)} \frac{ds}{s}$$

$$\lesssim N^{\frac{1}{2}} \int_0^\infty |\phi(s^{-1}) - \phi(\lambda)| (s\lambda e^{-(s\lambda-1)})^{N-1} \lambda ds.$$

Next we decompose the integral for $s\lambda \leq u_N$, $u_N < s\lambda < v_N$ and $s\lambda \geq v_N$, obtaining three terms I , II and III and where $u_N < 1$ and $v_N > 1$ will be suitably chosen later (around 1). For the first term, we have (since $x \rightarrow xe^{-(x-1)}$ is non-decreasing for $x \in (0, 1)$)

$$\begin{aligned} I &\lesssim N^{\frac{1}{2}} (u_N e^{-(u_N-1)})^{N-1} \lambda \|\phi\|_{L^\infty} \left(\int_{s\lambda \leq u_N} ds \right) \\ &\lesssim N^{\frac{1}{2}} (u_N e^{-(u_N-1)})^{N-1} \|\phi\|_{L^\infty}. \end{aligned}$$

For the third term; we similarly have (since $x \rightarrow xe^{-(x-1)}$ is decreasing for $x \in (1, \infty)$)

$$\begin{aligned} III &\lesssim N^{\frac{1}{2}} (v_N e^{-(v_N-1)})^{N-2} \left(\int_{s\lambda \geq v_N} (s\lambda) e^{-(s\lambda-1)} \lambda ds \right) \|\phi\|_{L^\infty} \\ &\lesssim N^{\frac{1}{2}} (v_N e^{-(v_N-1)})^{N-2} \|\phi\|_{L^\infty}. \end{aligned}$$

About the second term, using ω the uniform modulus of continuity of ϕ (and that $xe^{-(x-1)} \leq 1$ for every $x > 0$), it follows that

$$\begin{aligned} II &\lesssim N^{\frac{1}{2}} \int_{u_N \leq s\lambda \leq v_N} \omega\left(\frac{v_N - u_N}{s}\right) \lambda ds \\ &\lesssim N^{\frac{1}{2}} \omega(2\lambda(v_N - u_N)) |v_N - u_N|, \end{aligned}$$

where we used that u_N, v_N are around the value 1. Finally, we deduce that for every λ

$$\begin{aligned} |\mu_N(\lambda) - \phi(\lambda)| &\lesssim N^{\frac{1}{2}} (u_N e^{-(u_N-1)})^{N-1} \\ &\quad + N^{\frac{1}{2}} (v_N e^{-(v_N-1)})^{N-2} + N^{\frac{1}{2}} \omega(\lambda(v_N - u_N)) |v_N - u_N|. \end{aligned}$$

Now, let us write

$$u_N := 1 - \epsilon_N \quad \text{and} \quad v_N := 1 + \epsilon_N,$$

with $\epsilon_N \rightarrow 0$ as $N \rightarrow \infty$. We note that by a second order expansion,

$$\begin{aligned} \log \left(N^{\frac{1}{2}} (u_N e^{-(u_N-1)})^{N-1} \right) &= \frac{1}{2} \log(N) + (N-1) [\log(u_N) - (u_N - 1)] \\ &= \frac{1}{2} \log(N) - (N-1) \left[\frac{1}{2} (u_N - 1)^2 + \mathcal{O}(u_N - 1)^3 \right] \\ &= \frac{1}{2} \log(N) - (N-1) \left[\frac{1}{2} \epsilon_N^2 + \mathcal{O}(\epsilon_N^3) \right] \\ &\xrightarrow[N \rightarrow \infty]{} -\infty \end{aligned}$$

provided

$$(2.4) \quad \lim_{N \rightarrow \infty} \frac{N}{\log(N)} \epsilon_N^2 = \infty.$$

In this case,

$$\lim_{N \rightarrow \infty} N^{\frac{1}{2}} (u_N e^{-(u_N-1)})^{N-1} = 0.$$

Similarly, we have

$$\lim_{N \rightarrow \infty} N^{\frac{1}{2}} (v_N e^{-(v_N-1)})^{N-2} = 0,$$

and moreover since ω tends to 0 at 0 (with an order $\rho \in (0, 1]$: $\omega(s) \lesssim s^\rho$), we can choose ϵ_N such that

$$(2.5) \quad \lim_{N \rightarrow \infty} N^{\frac{1}{2}} |v_N - u_N| \omega(2\lambda(v_N - u_N)) \lesssim \lim_{N \rightarrow \infty} N^{\frac{1}{2}} \epsilon_N \omega(4\lambda \epsilon_N) = 0.$$

Indeed, take ϵ_N such that

$$N\epsilon_N^2 = \log(N)^2,$$

(which is possible for large enough integer N) then ϵ_N tends to 0 and (2.4) is satisfied. Moreover (2.5) follows from $\omega(s) \lesssim s^\rho$, due to the ρ -Hölder regularity of ϕ . For such ϵ_N , we finally conclude to

$$\lim_{N \rightarrow \infty} |\mu_N(\lambda) - \phi(\lambda)| = 0.$$

Then, using dominated convergence Theorem and then spectral theory in (2.3) implies (2.1). \square

Corollary 2.1. *Let $\delta \in (0, 1]$. For every smooth function ϕ and every L^2 -functions f, g , we have*

$$\begin{aligned} & \int_0^\infty \phi(\lambda) \langle dE_H(\lambda) f, g \rangle d\lambda \\ &= \lim_{N \rightarrow \infty} \frac{1}{(N-1)!} \int_0^\infty \phi(\lambda) \langle ((N-\delta)\lambda^{-1}H)^N e^{-\lambda^{-1}(N-\delta)H} f, g \rangle \frac{d\lambda}{\lambda} \\ &= \lim_{N \rightarrow \infty} \frac{1}{\Gamma(N+\delta)} \int_0^\infty \phi(\lambda) \langle (N\lambda^{-1}H)^{N+\delta} e^{-\lambda^{-1}NH} f, g \rangle \frac{d\lambda}{\lambda}. \end{aligned}$$

Proof. The case $\delta = 1$ is exactly the statement of Theorem 1.2. For $\delta \in (0, 1)$, we follow the same proof and replace $(N-1)! = \Gamma(N)$ by $\Gamma(N-\delta+1)$ (we recall that Stirling's formula remains valid for the Γ function, see (2.6) below). \square

Making an integration by parts in Theorem 1.2, we obtain a formula for the derivatives $\frac{d^k}{d\lambda^k} dE_H(\lambda)$ in terms of the semigroup. That is

Corollary 2.2. *The following equality holds in the weak sense: for an integer $k \geq 1$*

$$\frac{d^k}{d\lambda^k} dE_H(\lambda) = \lim_N \frac{1}{N!} \frac{d^k}{d\lambda^k} \left[\lambda^{-1} (N\lambda^{-1}H)^{N+1} e^{-N\lambda^{-1}H} \right]$$

for $\lambda > 0$.

Proof of Theorem 1.1. We first prove that 1) implies 2). Suppose that (1.1) is satisfied. For fixed N we have

$$\begin{aligned} \|H^N e^{-tH}\|_{L^p \rightarrow L^{p'}} &= \left\| \int_0^\infty \lambda^N e^{-t\lambda} dE_H(\lambda) \right\|_{L^p \rightarrow L^{p'}} \\ &\lesssim \int_0^\infty \lambda^N e^{-t\lambda} \lambda^{\frac{d}{2}(\frac{1}{p} - \frac{1}{p'})-1} d\lambda \\ &= \left(\int_0^\infty u^{N+\frac{d}{2}(\frac{1}{p} - \frac{1}{p'})-1} e^{-u} du \right) t^{-N-\frac{d}{2}(\frac{1}{p} - \frac{1}{p'})} \\ &= t^{-N-\frac{d}{2}(\frac{1}{p} - \frac{1}{p'})} \Gamma\left(N + \frac{d}{2}\left(\frac{1}{p} - \frac{1}{p'}\right)\right). \end{aligned}$$

Stirling's formula for the Gamma function (see [2, Appendix A.6])

$$(2.6) \quad \Gamma(x) \simeq x^{x-\frac{1}{2}} e^{-x} \sqrt{2\pi} \quad \text{for } x > 0,$$

shows that

$$\|H^N e^{-tH}\|_{L^p \rightarrow L^{p'}} \lesssim (N-1)! N^{\frac{d}{2}(\frac{1}{p} - \frac{1}{p'})} t^{-N-\frac{d}{2}(\frac{1}{p} - \frac{1}{p'})}.$$

This proves assertion 2).

We now prove that 2) implies 3). Suppose first that F is a Hölder continuous function with support in $[0, R]$. We apply Theorem 1.2 and obtain

$$\|F(H)\|_{L^p \rightarrow L^{p'}} \leq \limsup_N \frac{1}{(N-1)!} \int_{1/R}^\infty F(s^{-1}) \|((N-1)sH)^N e^{-s(N-1)H}\|_{p \rightarrow p'} \frac{ds}{s}$$

$$\begin{aligned}
&\lesssim \limsup_N \int_{1/R}^{\infty} F(s^{-1}) ((N-1)s)^N N^{\frac{d}{2}(\frac{1}{p}-\frac{1}{p'})} (s(N-1))^{-N-\frac{d}{2}(\frac{1}{p}-\frac{1}{p'})} \frac{ds}{s} \\
&\lesssim \int_{1/R}^{\infty} F(s^{-1}) s^{-\frac{d}{2}(\frac{1}{p}-\frac{1}{p'})} \frac{ds}{s} \\
&\leq R^{\frac{d}{2}(\frac{1}{p}-\frac{1}{p'})} \int_0^R F(s) \frac{ds}{s}.
\end{aligned}$$

We have proved that

$$(2.7) \quad \|F(H)\|_{L^p \rightarrow L^{p'}} \lesssim R^{\frac{d}{2}(\frac{1}{p}-\frac{1}{p'})} \int_0^R F(s) \frac{ds}{s}.$$

Next we extend the latter estimate to all bounded functions F with support in $[0, R]$. This can be achieved by classical approximation arguments. First assume that the support of F is contained in $[\eta, R]$ for some $\eta > 0$ and apply (2.7) to the convolution $F_\epsilon = \rho_\epsilon * F$ by a mollifier ρ_ϵ . We obtain

$$\|F_\epsilon(H)\|_{L^p \rightarrow L^{p'}} \lesssim (R + \epsilon)^{\frac{d}{2}(\frac{1}{p}-\frac{1}{p'})} \int_{\eta-\epsilon}^R F_\epsilon(s) \frac{ds}{s}.$$

Since $|F_\epsilon(s)| \leq \|F\|_\infty$ and the support of F_ϵ is contained in $[\eta/2, R + \eta/2]$ for $\epsilon < \eta/2$ one can apply the dominated convergence theorem to the RHS of the previous inequality. We obtain (2.7). Now for every bounded F with support in $[0, R]$ we can apply (2.7) to $\chi_{[\epsilon, R]} F$ and then let $\epsilon \rightarrow 0$. Assertion 3) is then proved.

Finally we prove that 3) implies 1). In order to do this, we fix $\lambda > 0$ and $\epsilon \in (0, \lambda)$, and apply 3) to $F(s) = \chi_{(\lambda-\epsilon, \lambda+\epsilon]}(s)$. It follows that

$$\begin{aligned}
\|\chi_{(\lambda-\epsilon, \lambda+\epsilon]}(H)\|_{L^p \rightarrow L^{p'}} &\lesssim (\lambda + \epsilon)^{\frac{d}{2}(\frac{1}{p}-\frac{1}{p'})} \int_{\lambda-\epsilon}^{\lambda+\epsilon} \frac{ds}{s} \\
&= (\lambda + \epsilon)^{\frac{d}{2}(\frac{1}{p}-\frac{1}{p'})} [\ln(\lambda + \epsilon) - \ln(\lambda - \epsilon)] \\
&\simeq (\lambda + \epsilon)^{\frac{d}{2}(\frac{1}{p}-\frac{1}{p'})} \frac{2\epsilon}{\lambda}.
\end{aligned}$$

Hence

$$\|\epsilon^{-1} \chi_{(\lambda-\epsilon, \lambda+\epsilon]}(H)\|_{L^p \rightarrow L^{p'}} \lesssim (\lambda + \epsilon)^{\frac{d}{2}(\frac{1}{p}-\frac{1}{p'})} \lambda^{-1}.$$

We let $\epsilon \rightarrow 0$ and obtain assertion 1). □

Remark 2.3.

- In the proof of 1) \Rightarrow 2) we can take $N = 0$ and obtain that 1) implies

$$\|e^{-tH}\|_{L^p \rightarrow L^{p'}} \lesssim t^{-\frac{d}{2}(\frac{1}{p}-\frac{1}{p'})}, \quad t > 0.$$

- Suppose that (1.1) holds. Let $\alpha > 0$ and apply assertion 3) with $F(s^\alpha)$ to obtain

$$\|F(H^\alpha)\|_{L^p \rightarrow L^{p'}} \lesssim R^{\frac{d}{2\alpha}(\frac{1}{p}-\frac{1}{p'})} \int_{\mathbb{R}} |F(s^\alpha)| \frac{ds}{s} = \alpha R^{\frac{d}{2\alpha}(\frac{1}{p}-\frac{1}{p'})} \int_{\mathbb{R}} |F(s)| \frac{ds}{s}.$$

We conclude by Theorem 1.1 that

$$\|dE_{H^\alpha}(\lambda)\|_{L^p \rightarrow L^{p'}} \leq C \lambda^{\frac{d}{2\alpha}(\frac{1}{p}-\frac{1}{p'})-1}, \quad \lambda > 0.$$

In particular, for $\alpha = \frac{1}{2}$

$$\|dE_{\sqrt{H}}(\lambda)\|_{L^p \rightarrow L^{p'}} \leq C \lambda^{d(\frac{1}{p}-\frac{1}{p'})-1}.$$

- Assume that the heat semigroup $(e^{-tH})_{t>0}$ satisfies the classical $L^p - L^2$ estimates $\|e^{-tH}\|_{L^p \rightarrow L^2} \lesssim t^{-\frac{d}{2}(\frac{1}{p}-\frac{1}{2})}$ for every $t > 0$ and some $p \in [1, 2]$. Then we observe that for every integer $N \geq 3$

$$\begin{aligned} \|H^N e^{-tH}\|_{L^p \rightarrow L^{p'}} &\leq \|e^{-\frac{t}{N}H}\|_{L^2 \rightarrow L^{p'}} \|H^N e^{-t(1-\frac{2}{N})H}\|_{L^2 \rightarrow L^2} \|e^{-\frac{t}{N}H}\|_{L^p \rightarrow L^2} \\ &\lesssim \left(\frac{t}{N}\right)^{-\frac{d}{2}(\frac{1}{p}-\frac{1}{p'})} \left(\frac{N}{t(1-\frac{2}{N})}\right)^N e^{-N} \\ &\lesssim t^{-N-\frac{d}{2}(\frac{1}{p}-\frac{1}{p'})} N^{\frac{d}{2}(\frac{1}{p}-\frac{1}{p'})} (Ne^{-1})^N \\ &\lesssim t^{-N-\frac{d}{2}(\frac{1}{p}-\frac{1}{p'})} N^{\frac{d}{2}(\frac{1}{p}-\frac{1}{p'})} (N-1)! \sqrt{N}, \end{aligned}$$

where we used Stirling's formula to obtain the last inequality. Therefore we see that the gap between this very general estimate with the one required in Theorem 1.1 is an extra term of order $N^{\frac{1}{2}}$.

3. RESTRICTION FROM DISPERSION

In this section we show that dispersive estimates for the semigroup generated by H imply restriction estimates and also $L^p - L^{p'}$ estimates for the derivatives $\frac{d^k}{d\lambda^k} dE_H(\lambda)$ up to some order. The result for the case $k = 0$ was already derived in [1] by a different proof. The result for $k \geq 1$ seems to be new.

Proposition 3.1. *Fix $1 \leq p < \frac{2d}{d+2} = 2_*$. Suppose that the semigroup $(e^{-zH})_{z \in \mathbb{C}^+}$ satisfies the following dispersive estimates:*

$$(3.1) \quad \|e^{-zH}\|_{L^p \rightarrow L^{p'}} \lesssim |z|^{-\frac{d}{2}(\frac{1}{p}-\frac{1}{p'})},$$

uniformly in z such that $\Re(z) > 0$. Then, for $\gamma > 0$

$$\|(NsH)^{N+\gamma} e^{-NsH}\|_{L^p \rightarrow L^{p'}} \lesssim s^{-\frac{d}{2}(\frac{1}{p}-\frac{1}{p'})} N^\gamma (N-1)!,$$

uniformly in $N \geq 1$ and $s > 0$ (the implicit constant only depends on $\gamma > 0$).

Remark 3.2. It follows from Cauchy's formula that (3.1) extends to the derivatives of the semigroup as follows: for $k \geq 1$ and every $z \in \mathbb{C}^+$

$$\|(\Re(z)H)^k e^{-zH}\|_{L^p \rightarrow L^{p'}} \lesssim |z|^{-\frac{n}{2}(\frac{1}{p}-\frac{1}{p'})}.$$

Proof. By the functional calculus, we have

$$(NsH)^{N+\gamma} e^{-NsH} = N! N^\gamma (sNH)^\gamma \int_{\Gamma_N} \frac{e^{-\zeta sH}}{(\zeta - N)^{N+1}} d\zeta,$$

where Γ_N is the circle of center N and of radius $N-1$. So we deduce that

$$\begin{aligned} \|(NsH)^{N+\gamma} e^{-NsH}\|_{L^p \rightarrow L^{p'}} &\leq N^{N+\gamma} N! \int_{\Gamma_N} \frac{\|(sH)^\gamma e^{-\zeta sH}\|_{L^p \rightarrow L^{p'}}}{|\zeta - N|^{N+1}} d\zeta \\ &\leq N^{N+\gamma} N! (N-1)^{-(N+1)} \int_{\Gamma_N} \|(sH)^\gamma e^{-\zeta sH}\|_{L^p \rightarrow L^{p'}} d\zeta \\ (3.2) \quad &\lesssim N! N^\gamma \int_0^{2\pi} \|(sH)^\gamma e^{-\zeta_\theta sH}\|_{L^p \rightarrow L^{p'}} d\theta, \end{aligned}$$

with $\zeta_\theta := N + (N-1)e^{i\theta}$. Writing (up to some numerical constant) with an integer $k \geq 1 + \gamma$

$$(sH)^\gamma e^{-\zeta sH} = s^\gamma \int_0^\infty (tH)^k e^{-(t+\zeta s)H} \frac{dt}{t^{1+\gamma}},$$

we deduce that (since $\gamma \in (0, k)$)

$$\begin{aligned}
\|(sH)^\gamma e^{-\zeta sH}\|_{L^p \rightarrow L^{p'}} &\lesssim s^\gamma \int_0^\infty \frac{t^k}{(t + s\Re(\zeta))^k (|t + s\zeta|)^{\frac{d}{2}(\frac{1}{p} - \frac{1}{p'})}} \frac{dt}{t^{1+\gamma}} \\
&\lesssim s^\gamma (s|\zeta|)^{-\frac{d}{2}(\frac{1}{p} - \frac{1}{p'})} \int_0^\infty \left(\frac{t}{t + s\Re(\zeta)} \right)^k \frac{dt}{t^{1+\gamma}} \\
&\lesssim (s|\zeta|)^{-\frac{d}{2}(\frac{1}{p} - \frac{1}{p'})} s^\gamma \left[\int_0^{s\Re(\zeta)} \left(\frac{t}{s\Re(\zeta)} \right)^k \frac{dt}{t^{1+\gamma}} + \int_{s\Re(\zeta)}^\infty \frac{dt}{t^{1+\gamma}} \right] \\
&\lesssim (s|\zeta|)^{-\frac{d}{2}(\frac{1}{p} - \frac{1}{p'})} \Re(\zeta)^{-\gamma} \\
&\lesssim (s|\zeta|)^{-\frac{d}{2}(\frac{1}{p} - \frac{1}{p'})}
\end{aligned}$$

where we used that $\Re(\zeta) \geq 1$ and $|t + s\zeta| \geq |s\zeta|$. Putting this estimate together with (3.2) yields with $\sigma := \frac{d}{2} \left(\frac{1}{p} - \frac{1}{p'} \right)$

$$\begin{aligned}
&\|(NsH)^{N+\gamma} e^{-NsH}\|_{L^p \rightarrow L^{p'}} \\
&\lesssim s^{-\sigma} N! N^\gamma \int_0^{2\pi} \left| N + (N-1)e^{i\theta} \right|^{-\sigma} d\theta \\
&\lesssim s^{-\sigma} N! N^\gamma \int_{-1}^1 ((N + (N-1)u)^2 + (N-1)^2(1-u^2))^{-\sigma/2} \frac{du}{\sqrt{1-u^2}} \\
&\lesssim s^{-\sigma} N! N^\gamma \left[\int_{-1}^0 (N^2(1+u)^2 + (N-1)^2(1+u))^{-\sigma/2} \frac{du}{\sqrt{1+u}} + \right. \\
&\quad \left. \int_0^1 N^{-\sigma} \frac{du}{\sqrt{1-u}} \right] \\
&\lesssim s^{-\sigma} N! N^\gamma \left[\int_0^{(N-1)^2} (1+v)^{-\sigma/2} \frac{dv}{N\sqrt{v}} + N^{-\sigma} \right] \\
&\lesssim s^{-\sigma} N! N^{\gamma-1} \simeq s^{-\sigma} (N-1)! N^\gamma,
\end{aligned}$$

where we used that $\sigma > 1$ since $p < 2_*$. \square

Corollary 3.3. *Assume that the semigroup $(e^{-zH})_{z \in \mathbb{C}^+}$ satisfies the dispersive estimate (3.1) for some $1 \leq p < \frac{2d}{d+2} = 2_*$. Then we have*

$$\|dE_H(\lambda)\|_{L^p \rightarrow L^{p'}} \lesssim \lambda^{\frac{d}{2}(\frac{1}{p} - \frac{1}{p'}) - 1}, \quad \lambda > 0.$$

In addition for an integer $k \leq d/2 - 1$, if $1 \leq p < \frac{2d}{d+2(k+1)}$ then

$$\left\| \frac{d^k}{d\lambda^k} dE_H(\lambda) \right\|_{L^p \rightarrow L^{p'}} \lesssim \lambda^{\frac{d}{2}(\frac{1}{p} - \frac{1}{p'}) - (k+1)}, \quad \lambda > 0.$$

Proof. The first assertion follows immediately from Proposition 3.1 and Theorem 1.1. For the second assertion we give for simplicity a proof for $k = 1$, the general case follows by iteration. By Corollary 2.2 we have in the weak sense

$$\begin{aligned}
\frac{d}{d\lambda} dE_H(\lambda) &= \lim_{N \rightarrow \infty} \frac{1}{N!} \frac{d}{d\lambda} \left[(N\lambda^{-1}H)^{N+1} e^{-\lambda^{-1}NH} \lambda^{-1} \right] \\
&= \lim_{N \rightarrow \infty} \frac{1}{N!} (NH)^{-1} \frac{d}{d\lambda} \left[(N\lambda^{-1}H)^{N+2} e^{-\lambda^{-1}NH} \right].
\end{aligned}$$

Following Proposition 3.1, we write

$$(N\lambda^{-1}H)^{N+2}e^{-N\lambda^{-1}H} = (N+2)!N^{N+2} \int_{\Gamma_N} \frac{e^{-\zeta\lambda^{-1}H}}{(\zeta-N)^{N+3}} d\zeta$$

and so

$$\frac{d}{d\lambda} \left[(N\lambda^{-1}H)^{N+2}e^{-N\lambda^{-1}H} \right] = (N+2)!N^{N+2}\lambda^{-2} \int_{\Gamma_N} \frac{\zeta H e^{-\zeta\lambda^{-1}H}}{(\zeta-N)^{N+1}} d\zeta.$$

Hence,

$$\frac{1}{N!}(NH)^{-1} \frac{d}{d\lambda} \left[(N\lambda^{-1}H)^{N+2}e^{-\lambda^{-1}NH} \right] = \frac{(N+2)!}{(N-1)!} N^N \lambda^{-2} \int_{\Gamma_N} \frac{\zeta e^{-\zeta\lambda^{-1}H}}{(\zeta-N)^{N+1}} d\zeta.$$

Using the dispersive estimate, it follows that

$$\begin{aligned} & \frac{1}{N!}(NH)^{-1} \left\| \frac{d}{d\lambda} \left[(N\lambda^{-1}H)^{N+2}e^{-\lambda^{-1}NH} \right] \right\|_{L^p \rightarrow L^{p'}} \\ & \lesssim \frac{(N+2)!}{(N-1)!} N^N \lambda^{-2} \int_{\Gamma_N} \frac{|\zeta| \|e^{-\zeta\lambda^{-1}H}\|_{L^p \rightarrow L^{p'}}}{|\zeta-N|^{N+1}} d\zeta \\ & \lesssim \frac{(N+2)!}{(N-1)!} \frac{N^N}{(N-1)^{N+1}} \lambda^{-2} \int_{\Gamma_N} |\zeta|^{1-\sigma} \lambda^\sigma d\zeta, \end{aligned}$$

with $\sigma := \frac{d}{2} \left(\frac{1}{p} - \frac{1}{p'} \right)$. Such integral was already computed in Proposition 3.1 and is uniformly bounded as soon as $1 - \sigma < -1$. This gives the desired estimate for $\frac{d}{d\lambda} E_H(\lambda)$. \square

REFERENCES

- [1] P. Chen, E.M. Ouhabaz, A. Sikora and L. Yan, Restriction estimates, sharp spectral multipliers and endpoint estimates for Bochner-Riesz means. Submitted (2011), <http://arxiv.org/abs/1202.4052>.
- [2] L. Grafakos, *Classical Fourier Analysis, Second Edition*. Graduate Texts in Mathematics **249**, Springer, New York, NY, 2009.
- [3] C. Guillarmou, A. Hassel and A. Sikora, Restriction and spectral multiplier theorems on asymptotically conic manifolds, to appear in *Analysis and PDE*, <http://arxiv.org/abs/1012.3780>.
- [4] L. Hörmander, *The analysis of linear partial differential operators*, **I, II**. Springer-Verlag, Berlin, 1983.

Frédéric Bernicot, CNRS - Université de Nantes, Laboratoire Jean Leray. 2, rue de la Houssinière, 44322 Nantes cedex 3. France,
Frederic.Bernicot@univ-nantes.fr

El Maati Ouhabaz, Institut de Mathématiques (IMB), Univ. Bordeaux, 351, cours de la Libération, 33405 Talence cedex, France,
Elmaati.Ouhabaz@math.u-bordeaux1.fr